

## EFFECT OF RECOMBINANT ERYTHROPOIETIN ON PERIPHERAL T LYMPHOCYTES

Kouichi Hisatomi, MD,<sup>a</sup> Masanobu Nakao, MD,<sup>b</sup> Tadashi Isomura, MD,<sup>a</sup> Kenichi Kosuga, MD,<sup>a</sup>  
and Kyogo Itoh, MD,<sup>b</sup> *Fukuoka, Japan*

Recombinant erythropoietin<sup>1</sup> has commonly been used to reduce the volume of homologous blood transfusion in patients undergoing cardiac operations in Japan. We<sup>2</sup> have reported that recombinant erythropoietin treatment not only increased levels of circulating erythrocytes but also improved indices of cell-mediated immunity, and we<sup>3</sup> have suggested that it might help to ameliorate or prevent the impairment of immune function that can occur after cardiac operations. We subsequently administered recombinant erythropoietin in a dosage of 200 U/kg per day for 7 days in a patient with postoperative erythroderma associated with pancytopenia after a cardiac operation. The level of interleukin-2 production was also found to increase in association with lessening of symptoms. In the present study, we investigated the effect of recombinant erythropoietin on peripheral T cell response with the use of a reverse-transcriptase polymerase chain reaction (PCR), which has currently been acknowledged to detect cytokines in vitro in immunologic studies. Peripheral blood mononuclear cells were obtained 7 days postoperatively from 10 patients who underwent valvular operations. Five of the patients without complications were given recombinant erythropoietin in a dosage of 100 U/kg for 7 days, and the other five patients were not. Samples were prepared by the Ficoll-Conray method. They were adjusted to a concentration of  $4 \times 10^6$  cells per milliliter and were incubated with 5, 10, 50, or 100 U/ml of recombinant erythropoietin for 6 hours at 37° C. Ribonucleic acid (RNA) was isolated by the RNeasy method (Qiagen Laboratories, Inc., Houston, Tex.). Cyclic deoxyribonucleic acid was prepared essentially as described by Becker, Quay, and Soukup.<sup>4</sup> The PCR primers used for this study were interleukin-2 primers

(sense primer: 5'-TGTACAGGATGCAACTCCTGTCTT; antisense primer: 5'-GTTAGTGTTGAGATGATGCTTTGAC). PCR was performed for 35 cycles. Reverse-transcriptase PCR-amplified messenger RNA was electrophoresed in a 1.5% agarose gel and stained with ethidium bromide.<sup>5</sup> T cells separated from peripheral blood mononuclear cells mixed with 5 or 10 U/ml of recombinant erythropoietin did not express interleukin-2, whereas the samples mixed with 50 or 100 U/ml of recombinant erythropoietin had decreased expression regardless of whether patients received recombinant erythropoietin for 7 days after the operation. These findings suggest that, at the messenger RNA level, recombinant erythropoietin may inhibit production of interleukin-2 from T cells in proportion to the dose. These results suggest that further study will be needed to reveal the true effect of recombinant erythropoietin on cell-mediated immune responses.

### REFERENCES

1. Jacobs K, Shoemaker C, Rudersdorf R, et al. Isolation and characterization of genomic and cDNA clones of human erythropoietin. *Nature* 1985;313:806-10.
2. Hisatomi K, Isomura T, Kawara T, et al. Changes in lymphocyte subsets, mitogen responsiveness, and interleukin-2 production after cardiac operations. *J THORAC CARDIOVASC SURG* 1989;98:580-91.
3. Hisatomi K, Isomura T, Galli SJ, Yasunaga H, Hayashida N, Ohishi K. Augmentation of interleukin-2 production after cardiac operations in patients treated with erythropoietin. *J THORAC CARDIOVASC SURG* 1992;104:278-83.
4. Becker S, Quay J, Soukup J. Cytokine (tumor necrosis factor, IL-6 and IL8) production by respiratory syncytial virus-infected human alveolar macrophages. *J Immunol* 1991;147:4307-12.
5. Hoshino T, Yamada A, Honda J, et al. Tissue-specific distribution and age-dependent increase of human CD11b+ T cells. *J Immunol* 1993;151:2237-46.

From the Second Department of Surgery<sup>a</sup> and the Department of Immunology,<sup>b</sup> Kurume University School of Medicine, Fukuoka, Japan.

*J THORAC CARDIOVASC SURG* 1995;109:809

Copyright © 1995 by Mosby-Year Book, Inc.

0022-5223/95 \$3.00 + 0 12/8/57781